

HELPING CHILDREN AFTER A DISASTER

A catastrophe such as an earthquake, hurricane, tornado, fire, flood, or violent acts is frightening to children and adults alike. It is important to **acknowledge the frightening parts of the disaster when talking with a child** about it. Falsely minimizing the danger will not end a child's concerns. Several factors affect a child's response to a disaster:

- **The way children see and understand their parents' responses are very important.** Children are aware of their parents' worries most of the time, but they are particularly sensitive during a crisis. Parents should admit their concerns to their children, and also stress their abilities to cope with the situation.
- **A child's reaction also depends on how much destruction and/or death he or she sees** during and after the disaster. If a friend or family member has been killed or seriously injured, or if the child's school or home has been severely damaged, there is a greater chance that the child will experience difficulties.
- **A child's age affects how the child will respond to the disaster.** For example, six-year olds may show their worries about a catastrophe by refusing to attend school, whereas adolescents may minimize their concerns, but argue more with parents and show a decline in school performance. It is important to explain the event in words the child can understand.

Following a disaster, people may develop Posttraumatic Stress Disorder (PTSD), which is psychological damage that can result from experiencing, witnessing, or participating in an overwhelmingly traumatic (frightening) event. Children with this disorder have repeated episodes in which they **re-experience the traumatic event**. Children **often relive the trauma through repetitive play**. In young children, **upsetting dreams** of the traumatic event may change into **nightmares** of monsters, of rescuing others, or of **threats to self or others**. PTSD rarely appears during the trauma itself. Though its symptoms can occur soon after the event, the disorder often surfaces several months or even years later.

Parents should be alert to these changes in a child's behavior:

- **Refusal to return to school and "clinging" behavior**, including shadowing the mother or father around the house
- **Persistent fears** related to the catastrophe (such as fears about being permanently separated from parents)
- **Sleep disturbances** such as nightmares, screaming during sleep and bedwetting, persisting more than several days after the event
- **Loss of concentration and irritability**
- **Startled easily; jumpy**
- **Behavior problems**, for example, misbehaving in school or at home in ways that are not typical for the child
- **Physical complaints** (stomachaches, headaches, dizziness) for which a physical cause cannot be found
- **Withdrawal from family and friends**, sadness, listlessness, **decreased activity**, and preoccupation with the events of the disaster

Professional advice or treatment for children affected by a disaster--especially those who have witnessed destruction, injury or death--can help prevent or minimize PTSD. For more information, see Facts for Families (<http://www.aacap.org/publications/pubcat/facts.htm>). Facts for Families© is developed and distributed by the American Academy of Child and Adolescent Psychiatry (AACAP).

Distributed by: Directorate of Health Promotion and Wellness
U.S. Army Center for Health Promotion and Preventive Medicine
Aberdeen Proving Ground, MD 21010-5403
Commercial 410-436-4656 or DSN 584-4656
<Http://chppm-www.apgea.army.mil/dhpw>